



Dermatology	' Order	Form
Fax completed form to: _		

		PATIENT	ΓINFORMATIO	N		
Patient Name:		Date of Birth:			Referral Date:	
Address:				City/State/Zip	):	
Home Phone:		Cell Phone:		İ	Work Phone:	
Secondary Contact:		Height:	Weight:		Male Female	
Patient Diagnosis & ICD	D-10:					
Allergies:						
PROVIDER INFORMATION						
Physician Name:		Lic.#:		DEA #:		
Practice Name:				NPI#:		
Address:				City/State/Zip	):	
Office Contact:		Phone:			Fax:	
Supervisory Physician (i	if applicable):					
		PLE	ASE ATTACH			
Patient demographi	ics & front/back copy of all insurance cards (pres	scription & medical)	TB lab results within la	st 12 months <i>(Ste</i>	lara, Simponi Aria, Ilumya & Infliximabs only,	)
Recent office visit no	otes, history & physical, lab & pertinent proced	ure results	HBV lab results within	last 12 months <i>(li</i>	nfliximabs & Simponi Aria only)	
Current medication	list & list of prior medications tried and failed (v	with dates)	Letter of medical neces	sity if drug dosin	g or indication is outside of FDA guidelines	
		NURSIN	G & LAB ORDEI	RS		
Nurse Orders: Nurse to	provide assessment, teaching, lab draws, med	lication administration an	d vascular access device ins	ertion and/or ma	nagement per physician orders.	
	% - 5-10mL flush pre and post infusion and as				flush after post-infusion NS flush if indicate	d to maintain line
Lab Orders:		•	Lab Date & Frequency		·	
		PRESCR	IPTION ORDER	S		
Anaphylaxis Kit:	Epinephrine 0.3mg IM as needed		rtef 250mg-500mg IV as ne		Solu-Medrol 60mg - 125m	na IV as needed
(Check all that apply)			ation 500 ml IV over 30 min		Other	ig iv as inceaca
Pre-Medications:	Acetaminophenmg PO	minutes prior to				
(Check all that apply)	Diphenhydraminemg	POOR IV infus		or to infusion	Other	
	plies for vascular access line care, drug adminis				other	
PRODUCT			'ION INFORMA'			REFILLS
Is this a first dose?	Yes No If No, when was last dose given?		When is patient due for nex	rt dose?		
ILUMYA	100mg SC injection at 0 and 4 weeks then eve	ery 12 weeks				
INFLIXIMAB	Induction:mg/kg or	mg IV infusion via	gravity OR pump	over at least 2 h	ours at weeks 0, 2, and 6	NONE
Avsola	Maintenance:mg/kg	mg IV infusion via	gravity OR pum	o over at least 2 h	ours every weeks	
Inflectra	(Note: Round to nearest 100mg for Medicaid p		3. 7			
Remicade	If Remicade infusion tolerated, adjust infusio		facturer nackage insert			
Renflexis						
SIMPONI ARIA	2 mg/kg IV infusion via gravityOR		es at weeks 0 and 4, and ev			
SPEVIG0		Additional 900 mg IV infus	sion over 90 minutes one we	eek after initial do	se if flare symptoms persist	
	Psoriasis Adult Subcutaneous		. (       11 45	12		
	For patients <= 100 kg, 45 mg SC injection For patients > 100 kg, 90 mg SC injection					
	Psoriasis Pediatric Patients 6 to 17 (base			Z WCCK3		
CTEL ADA	For patients <= 60 kg, 0.75 mg/kg SC inju			5		
STELARA	For patients 60 kg — 100kg, 45 mg SC injection initially and 4 weeks later, then every 12 weeks					
	For patients >100kg, 90 mg SC injection i	nitially and 4 weeks later,	then every 12 weeks			
	Psoriatic Arthritis Adult					
	45 mg SC injection initially and 4 weeks la	ater, followed by 45 mg SC				
	For nationts with so existent moderate to	o covoro plaguo peoriasia.	woighing > 100 kg 00 cc = 0	Cininction initial	ly and 1 wooks later than avery 12 waster	
VOLAID	For patients with co-existent moderate-to		weighing >100 kg, 90 mg S	C injection initial	ly and 4 weeks later, then every 12 weeks	
XOLAIR	150 or 300 mg SC injection once every	y 4 weeks		C injection initial	ly and 4 weeks later, then every 12 weeks	
IG	· ·	y 4 weeks		C injection initial	ly and 4 weeks later, then every 12 weeks	
IG OTHER	150 or 300 mg SC injection once every	y 4 weeks <b>to Immunoglobulin For</b>	m			nce companies.

Prescriber's Signature **Dispense as Written** 

**Print Name** 

Date

Prescriber's Signature **Substitution Permitted**  **Print Name** 

Date





