## Gastroenterology Order Form





Fax complete	d form to:	specialty infusion services	infusion solution
	PATIENT INFOR	MATION	
Patient Name:	Date of Birth:	Referral Date:	
Address:	- Date of Strain	City/State/Zip:	
Home Phone:	Cell Phone:	Work Phone:	
Secondary Contact:	Height: Weight:	Male Female	
Patient Diagnosis & ICI			
Allergies:			
	PROVIDER INFO	RMATION	
Physician Name:	Lic.#:	DEA #:	
Practice Name:	•	NPI#:	
Address:		City/State/Zip:	
Office Contact:	Phone:	Fax:	
Supervisory Physician	(if applicable):		
	PLEASE ATT	ACH	
Current medication Line access docum  Nurse Orders: Nurse to	It ist & list of prior medications tried and failed (with dates) HBV lab resentation/verification if applicable  NURSING & LAB to provide assessment, teaching, lab draws, medication administration and vascular accee 100% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL01	<u> </u>	ers.
	PRESCRIPTION	ORDERS	
Anaphylaxis Kit: (Check all that apply)		/ over 30 minutes as needed Other	rol 60mg - 125mg IV as needed
Pre-Medications:	Acetaminophenmg POminutes prior to infusion  Diphenhydraminemg PO <b>OR</b> IVminutes	Solu-Medrolmg IVminutes prior to infl s prior to infusion Other	usion
(Check all that apply)	Diphenhydramine mg PO <b>OR</b> IV minutes pplies for vascular access line care, drug administration kit(s), pump, and IV pole will be p		
PRODUCT	PRESCRIPTION INFO	, , , , , , , , , , , , , , , , , , ,	REFILLS
			REFILLS
Is this a first dose?		nt due for next dose?	
ENTYVIO	Induction: 300mg IV infusion via gravityOR pump over 30 minute Maintenance: 300mg IV infusion via gravityOR pump over 30 minute gravity	es at week 0, 2, and 6 nutes every weeks	NONE
INFLIXIMAB	Induction:mg/kg ormg IV infusion via gravityOR-	pump over at least 2 hours at weeks 0, 2, and 6	NONE
Avsola	Induction:mg/kg ormg IV infusion via gravityOR  Maintenance: mg/kg mg/V infusion via gravityOR		INUNE
Inflectra	(Note: Round to nearest 100mg for Medicaid patients)	P pump over at least 2 hours every weeks	
Remicade			
5 0 .	If Remicade infusion tolerated, adjust infusion time according to manufacturer package	ge insert.	

By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

weeks after induction and every

Prescriber's Signature Print Name Date
Dispense as Written

Maintenance: 90mg SubQ injection

Prescriber's Signature Substitution Permitted **Print Name** 

pump over at least 1 hour x 1 dose

pump over at least 1 hour x 1 dose

weeks thereafter

Date

NONE

NONE







Renflexis

STELARA

OTHER

For patients 55kg or less administer 260mg IV infusion via gravity --- OR--- pump over at least 1 hour x 1 dose

Induction (Adult Dosing -Based on body weight of patient at time of dosing):

For patients more than 85kg administer 520mg IV infusion via gravity --- OR---

For patients more than 55kg to 85kg administer 390mg IV infusion via gravity --- OR---