

Neurology Order Form



Fax completed form to: _____

| PATIENT INFORMATION | | | |
|--|--|--|--|
| Patient Name: | Date of Birth: | Referral Date: | |
| Address: | | City/State/Zip: | |
| Home Phone: | Cell Phone: | Work Phone: | |
| Secondary Contact: | Height: | Weight: | Male Female |
| Patient Diagnosis & ICD-10: | | | |
| Allergies: | | | |
| PROVIDER INFORMATION | | | |
| Physician Name: | Lic.#: | DEA #: | |
| Practice Name: | | NPI#: | |
| Address: | | City/State/Zip: | |
| Office Contact: | Phone: | Fax: | |
| Supervisory Physician (if applicable): | | | |
| PLEASE ATTACH | | | |
| Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicable Quantitative serum Immunoglobulin lab results (<i>Uplizna only</i>) TB lab results within last 12 months (<i>Uplizna only</i>) | | Vaccine status (any vaccination) and documentation of any recent vaccinations HBV lab results within last 12 months (<i>Uplizna only</i>) Date of diagnosis, current FVC%, ALSFRS-R score, and JourneyMate form (<i>Radicava only</i>) Anti-acetylcholine receptor (AChR) antibody positive results (<i>Vyvgart</i>) Letter of medical necessity if drug dosing or indication is outside of FDA guidelines | |
| NURSING & LAB ORDERS | | | |
| Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL ---OR--- 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders: Lab Date & Frequency: | | | |
| PRESCRIPTION ORDERS | | | |
| Anaphylaxis Kit: (Check all that apply) | Epinephrine 0.3mg IM as needed Diphenhydramine _____ mg IV as needed | Solu-cortef 250mg-500mg IV as needed NS Hydration 500 ml IV over 30 minutes as needed | Solu-Medrol 60mg - 125mg IV as needed Other _____ |
| Pre-Medications: (Check all that apply) | Acetaminophen _____ mg PO _____ minutes prior to infusion Diphenhydramine _____ mg PO ---OR--- IV _____ minutes prior to infusion | Solu-Medrol _____ mg IV _____ minutes prior to infusion | Other _____ |
| Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary | | | |
| PRODUCT | PRESCRIPTION INFORMATION | | REFILLS |
| Is this a first dose? Yes No | If No, when was last dose given? _____ When is patient due for next dose? _____ | | |
| RADICAVA | Induction: 60mg IV infusion via gravity ---OR--- pump over 1 hour daily for 14 days followed by 14 day drug-free period Maintenance: 60mg IV infusion via gravity ---OR--- pump over 1 hour daily for 10 days out of 14 day period followed by 14 day drug-free periods | | NONE |
| UPLIZNA | Induction: 300mg IV infusion via gravity ---OR--- pump over approximately 90 minutes at 0 and 2 weeks and CBC lab testing every _____ months Maintenance: (starting 6 months from first infusion) 300mg IV infusion via gravity ---OR--- pump over approximately 90 minutes every 6 months | | NONE |
| VYEPTI | 100mg IV infusion via gravity ---OR--- pump over approximately 30 minutes every 12 weeks 300mg IV infusion via gravity ---OR--- pump over approximately 30 minutes every 12 weeks | | |
| VYVGART | 10mg/kg IV infusion via gravity ---OR--- ---OR--- pump over at least 1 hour once every week for 4 weeks *Up to max of 1200mg for patient weight of 120kg+ (Total volume is 125ml in NS solution) Administer additional treatment cycles every 50 days ---OR--- Prescriber to evaluate treatment cycle frequency after completion of initial treatment cycle According to the Package Insert: Administer subsequent treatment cycles based on clinical evaluation; the safety of initiating subsequent cycles sooner than 50 days from the start of the previous treatment cycle has not been established. | | |
| VYVGART HYTRULO | 1,008mg/11,200 units subcutaneous injection over approximately 30 to 90 seconds in cycles of once weekly injections for 4 weeks Administer additional treatment cycles every 50 days ---OR--- Prescriber to evaluate treatment cycle frequency after completion of initial treatment cycle According to the Package Insert: Administer subsequent treatment cycles based on clinical evaluation; the safety of initiating subsequent cycles sooner than 50 days from the start of the previous treatment cycle has not been established. | | |
| IG | Refer to Immunoglobulin Form | | |
| SOLIRIS/ULTOMIRIS | Refer to Soliris or Ultomiris Order Form | | |
| OTHER | | | NONE |
| By signing this form and utilizing our services, you are authorizing Mosaic to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies. | | | |

Prescriber's Signature _____ Print Name _____ Date _____
 Dispense as Written

Prescriber's Signature _____ Print Name _____ Date _____
 Substitution Permitted