Ultomiris Order Form





Fax completed form to:

| PATIENT INFORMATION | | | | | | |
|---|---|-------------------|--|------------------------------|---------|--|
| Patient Name: Date of Birth: | | | Referral Date: | | | |
| Address: | | | City/State/Zip: | | | |
| Home Phone: | | | | Work Phone: | | |
| Secondary Contact: Height: | | Weight: | Male Female | | | |
| Patient Diagnosis & ICD-10: | | | | | | |
| Allergies: PROVIDER INFORMATION | | | | | | |
| | | | | | | |
| Physician Name: Practice Name: | LIC.#. | | NPI#: | | | |
| Address: | | | City/State/Zip: | | | |
| Office Contact: Phone: | | | Fax: | | | |
| Supervisory Physician (if applicable): | | | | | | |
| PLEASE ATTACH | | | | | | |
| Patient demographics & front/back copy of all insurance cards (prescription & medical) Vaccine status (any vaccination) and documentation of any recent vaccinations | | | | | | |
| Recent office visit notes, history & physical, lab & pertinent procedure results Clinical documentation on any recent meningococcal infections | | | | | | |
| Current medication list & list of prior medications tried and failed (with dates) Documentation of a meningococcal vaccination | | | | | | |
| Line access documentation/verification if applicable Letter of medical necessity if drug dosing or indication is outside of FDA guidelines | | | | | | |
| NURSING & LAB ORDERS | | | | | | |
| | | | | | | |
| Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 10units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line | | | | | | |
| Lab Orders: Lab Orders: | | | | | | |
| | | | | | | |
| PRESCRIPTION ORDERS | | | | | | |
| Anaphylaxis Kit: | Epinephrine 0.3mg IM as neededSolu-cortef 250mg-500mg IV infusion as neededSolu-Medrol 60mg - 125mg IV infusion as needed | | | | | |
| (Check all that apply) Diphenhydramine mg IV infusion as needed NS Hydration 500 ml IV infusion over 30 minutes as needed Other | | | | | | |
| Pre-Medications: Acetaminophenmg P0minutes prior to infusion Solu-Medrolmg IV infusionminutes prior to infusion | | | | | | |
| (Check all that apply) Diphenhydramine mg POOR IV infusionminutes prior to infusion Other | | | | | | |
| Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary | | | | | | |
| | | | | | DEFILLO | |
| | | | | | | |
| Is this a first dose? Yes No If No, when was last dose given? When is patient due for next dose? | | | | | | |
| Is the prescriber enrolled in the Ultomiris REMS program? Yes No | | | | | | |
| Ultomiris | Loading Dose | | | | | |
| PNH and aHUS | For patients 5-10kg administer 600mg IV infusion via | gravity 0R | | | | |
| | For patients 10-20kg administer 600mg IV infusion via | gravity 0R | DR pump over at least 0.8 hours | | NONE | |
| | For patients 20-30kg administer 900mg IV infusion via | gravity 0R | pump over at least 0.6 hours | | | |
| | For patients 30-40kg administer 1,200mg IV infusion via gravity OR pump over at least 0.5 hours | | NONL | | | |
| PNH, aHUS and gMG | For patients 40-60kg administer 2,400mg IV infusion via | | | | | |
| | For patients 60-100kg administer 2,700mg IV infusion v | | | pump over at least 0.6 hours | | |
| | For patients >100kg administer 3,000mg IV infusion via gravityOR pump over at least 0.4 hours | | | | | |
| PNH and aHUS | Maintenance Dose | | | | | |
| | For patients 5-10kg administer 300mg IV infusion via | gravityOR | pump over at least 0.8 hours every 4 weeks | | | |
| | For patients 10-20kg administer 600mg IV infusion via | gravityOR | pump over at least 0.8 hours every 4 weeks | | | |
| | For patients 20-30kg administer 2,100 IV infusion via gravity OR pump over at least 1.3 hours every 8 weeks | | | | | |
| PNH, aHUS and gMG | For patients 30-40kg administer 2,700mg IV infusion via | | | | | |
| | For patients 40-60kg administer 3,000mg IV infusion via | | | • | | |
| | For patients 60-100kg administer 3,300mg IV infusion v | | | | | |
| | For patients >100kg administer 3,600mg IV infusion via | gravityOR | | | | |
| OTHER | | | | | NONE | |
| By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies. | | | | | | |
| | | | | | | |

Prescriber's Signature <u>Dispense as Written</u> Print Name

Date

Prescriber's Signature Substitution Permitted Print Name

Date

©2022 Amerita Inc. All rights reserved. AME MOS Ultomiris ORDER FORM 06-17-22

mosaiciv.com



